



ISSN: 0976-3376

Available Online at <http://www.journalajst.com>

ASIAN JOURNAL OF
SCIENCE AND TECHNOLOGY

Asian Journal of Science and Technology
Vol. 12, Issue, 02, pp.11553-11560, February, 2021

RESEARCH ARTICLE

CARCASS AND NON-CARCASS YIELD CHARACTERISTICS OF HORRO SHEEP FED ELEPHANT GRASS (*PENNISETUM PURPUREUM*) ENSILED WITH DIFFERENT PROPORTION OF DOLICHOS LABLAB (*LABLAB PURPUREUS*)

Tusa Gemechu^{1*}, Meseret Girma² and Mitiku Eshetu²

¹Oromia Agricultural Research Institute, Bako Agricultural Research Center, Bako, Ethiopia.

²Haramaya University School of Animal and Range Science, Haramaya, Ethiopia.

ARTICLE INFO

Article History:

Received 27th November, 2020

Received in revised form

19th December, 2020

Accepted 14th January, 2021

Published online 28th February, 2021

Key words:

Carcass, Dolichos Lablab,
Elephant Grass, Horro Sheep,
Non-Carcass, Silage.

ABSTRACT

The study was conducted at Bako Agricultural Research Center with the aim to evaluate the effects of Elephant grass ensiled with different proportion of Dolichos lablab on carcass and non-carcass yield. Elephant grass was harvested after eight weeks of regrowth. While, Dolichos lablab harvested at 50% blooming stage and they were chopped separately, mixed, added with 3% molasses and ensiled. A total of twenty intact male yearling Horro sheep were stratified into four groups based on initial body weight in RCBD and assigned to the dietary treatments randomly. Dietary treatments were arranged as T1, T2, T3 and T4 respectively for Elephant grass ensiled alone, with 10, 20 and 30% of Dolichos lablab. Lambs in all treatments were supplemented with wheat bran at 1.5% body weight. The study comprised 90 days of feeding trial for evaluation of carcass parameters. Increasing the inclusion levels of Dolichos lablab in the Elephant grass silage significantly increased slaughter weight ($P<0.01$), empty body weight, hot carcass weight ($P<0.001$), Hind and fore quarter, Dressing percentage on SW basis, EBW basis and REMA ($P<0.05$). The dietary treatments were significantly influenced ($P<0.05$) mean weight of tail, heart, liver with gall bladder, omasum-abomasum, large and small intestine, head with tongue and total edible offal components. Feeding of ensiled Elephant grass with different proportion of Dolichos lablab significantly affect weight values of Skin, gut contents and head without tongue of Horro lambs. Whereas, the rest parameters of non-edible offal components were not influenced. Therefore, addition of 20-30% Dolichos lablab in Elephant grass silage enhanced the nutritive value of Elephant grass silage for better growth and carcass yields of Horro sheep.

Citation: Tusa Gemechu, Meseret Girma and Mitiku Eshetu. 2021. "Carcass and Non-carcass Yield Characteristics of Horro Sheep Fed Elephant Grass (*Pennisetum purpureum*) Ensiled with Different Proportion of Dolichos Lablab (*Lablab purpureus*)", *Asian Journal of Science and Technology*, 12, (02), 11553-11560.

Copyright © 2021, Tusa Gemechu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Small ruminant production is a major component of the livestock sector in Ethiopia. According to CSA (2018) report, the total small ruminant population in Ethiopia is estimated to be about 64.04 million, out of which 31.3 million are sheep. At smallholder level, sheep are the major source of food security serving a diverse function including cash income, savings, cash for fertilizer purchase and sociocultural functions (Solomon et al., 2013). However, the productivity of sheep per head is low mainly because of inadequate year round nutrition, both in terms of quantity and quality, unimproved genetic potential and due to prevalence of diseases and parasites (Markos et al., 2006). Nowadays most of natural pasturelands are put under intensified crop production due to the increasing human population pressure. The remaining land is heavily grazed and cannot meet the nutritional requirement of livestock, resulting in reduced growth rate, low production,

poor fertility and high mortality particularly in the dry season (Solomon, 2004).

Aftermath grazing and crop residue accounts for 60 to 70% of available basal diet in the highlands of Ethiopia. Such feeds are inherently low in nutritive value such as protein, digestible energy and minerals, which may result in sub-optimal rumen fermentation and lowered animal performance (Malede and Tekele, 2014). As a result, animals could not realize their genetic potential within the growth period for the breed. To solve this problem, there are options like supplementing animals with agro-industrial by-products such as different oil seed cakes and brans from edible oil and flour processing industries. However, they are costly and not easily available everywhere. The other option of increasing livestock production can be achieved through cultivation of high quality forages with high yielding ability that are adapted to different environmental stresses (Muia et al., 2001; Tessema et al., 2010). Improved pasture crops have many advantages such as providing quality feed and improving soil fertility. The cost

incurred for perennial pasture production is high only during the establishment year (J rgensen *et al.*, 2010) but declined thereafter which could serve for the way forward in feed resources improvement strategies in Ethiopia at large. Amongst the promising forage species promoted in Ethiopia, Elephant grass could play an important role in providing a significant amount of high-quality forage to the livestock industry (Ndikumana, 1996; Tessema, 2005) both under the smallholder farmer and intensive livestock production systems with appropriate management practices (Alemayehu, 2004). Uses of improved forage and tree legumes as supplementary options for livestock have been investigated in Ethiopia and elsewhere in Africa (Solomon *et al.*, 2004; Ajebu *et al.*, 2008 and Adegun, 2014). Improved forages could be fed to the animals by direct grazing, cut and carry, or stored and conserved as hay or silage. The potential of legumes for grazing is limited due to their susceptibility to trampling and the preferential grazing by livestock that adds pressure to legumes (Phelan *et al.*, 2015). Additionally, fresh feeding of legumes is limited due to the seasonality of rain which reduces the independence from protein-rich feedstuffs during the dry season. Moreover, legumes feeding as hay restricted by high dry-matter losses occurring during drying, transport and storage furthermore, because of the practical implications of season and plant physiology that may prevent a timely harvest/drying at the optimal time for hay making (Titterton and Bareeba, 1999).

The limitations due to leaf shattering, harvest timing and availability in the dry season are minimized when forages are conserved as silage, making this probably the most suitable option to feed tropical legumes to ruminants. The development of small-scale intensive systems and the rising costs of concentrate feeds, made the conservation of forage crops an integral part of livestock production (Mapiye *et al.*, 2007). Conserving high quality forages reduces the need to purchase concentrates for use in ruminant rations. Surplus and cultivated quality forages should be conserved during the wet season for use during the dry season. To this effect, silage-making is a common means of preserving surplus forage which could be fed to livestock during periods of scarcity. By conserving excess forage produced during the wet season to silage, the low production and productivity of livestock during the dry season due to scarcity of forage can be ameliorated (Wong, 2000). Conservation of cereals/grasses produces silages of high energy but low protein concentration (Mhere *et al.*, 2002). On the other hand, legumes alone do not ensile well owing to their high moisture content and high buffering capacity, resulting in high effluent losses and unstable silage of high pH (Pahlow *et al.*, 2002). Incorporation of legumes in cereal silage increases protein concentration improves silage intake and its nutritional quality thereby enhances ruminant animal performance (Ngongoni *et al.*, 2008). A number of Elephant grass and Dolichos lablab varieties are proved to produce high dry matter yield and adapted to a wide range of agro ecologies across the country. However, they contribute little to the much needed improvement of livestock production because data regarding the conservation of mixed grass-legume forages as silage was scarce. Similarly, feeding of silage in general and grass-legume mixed silage in particular for sheep as a basal diet is not practiced in the study area. Therefore, an alternative feed resources mainly conserved grass-legume mixed silage to enhance the productivity of small ruminants and determining the nutrient concentration of

Elephant grass ensiled alone and with different proportion of Dolichos lablab is yet not investigated and exploited. Thus, the study was undertaken with the objective to evaluate the effects of Elephant grass ensiled with different proportion of Dolichos lablab on carcass and non-carcass yield of Horro sheep.

MATERIALS AND METHODS

Description of the Study Area: The experiment was conducted at Bako Agricultural Research Center which is located in Oromia Regional State, West Shoa Zone at about 257 Km from the capital city Addis Ababa on the way to Nekemte town.

Forage Production and Preservation: Pre-existing Elephant grass accession (ILRI 14984) was cut at about 50cm height above the ground in order to re-growth and later for silage making. In other fields, Dolichos lablab seeds was sown at recommended rate. The cutting of Elephant grass was mainly to match its appropriate harvesting time with Dolichos lablab during silage making. Elephant grass was harvested after eight weeks of re-growth, while the Dolichos Lablab was harvested at 50% blooming stage.

The Elephant grass and Dolichos Lablab were separately chopped in to smaller pieces using machine chopper driven by tractor before ensiling. Before ensiling plant materials were mixed according to treatment combinations. To improve fermentation quality of ensiling materials 3% sugar cane molasses was sprinkled uniformly to all treatments over the chopped fodders and thoroughly mixed. For the ease of application, the sugar cane molasses was diluted with water 3:1 ratio before use. The chopped and mixed fodders ready for ensiling were preserved in separate pit silos lined with polythene sheets. It was compacted by trampling with feet, covered with polythene sheets and heap the soil on the pit to maintain anaerobic conditions during storage. The silages was opened after 45 days and used in a feeding trial to evaluate carcass and non-carcass of Horro sheep fed ad libitum Elephant grass and Dolichos lablab mixed silage.

Experimental Animals and Management: Before the commencement of actual experiment the experimental house, pens, feeding and watering trough were properly maintained, cleaned and disinfected. A total of twenty intact male yearling Horro sheep with initial body weight of 19.47 ± 1.76 Kg (mean \pm SD) used for this experiment were bought from local market. Age of the animals were determined by their dentition and information obtained from owners. The experimental animals were ear-tagged for identification, quarantined for fifteen days and vaccinated against common infectious diseases in the area. Thereafter, the experimental animals were assigned into different treatments after which the animals were randomly put in to a separate well aerated pen having a feed trough.

Experimental Design and Treatments: A randomized complete block design (RCBD) were used. The experimental animals were grouped into five blocks of four animals each based on their initial body weight. Animals in a block were assigned to one of the four treatments and uniformly supplemented with wheat bran 1.5% of their body weight across all the treatments.

The arrangement of dietary treatments were as follows;

- T1: Elephant grass ensiled alone
- T2: Elephant grass ensiled with 10% Dolichos Lablab
- T3: Elephant grass ensiled with 20% Dolichos Lablab
- T4: Elephant grass ensiled with 30% Dolichos Lablab

Carcass Analysis: All the experimental animals were slaughtered by severing the jugular vein using sharp knives after an overnight fasting. During slaughtering, the blood was drained and weighed. Weight of visceral organs like kidneys, liver with gall bladder, lungs with trachea, kidney fat, abdominal fat, omental fat, heart, spleen, genital organs, the entire alimentary canal (esophagus, reticulo-rumen, omasum-abomasum, small intestine and large intestine), full gut, empty gut, tail, tongue and head was recorded separately. The weight of the gut contents were measured by differences of full gut contents and empty gut contents. Empty body weight was determined by subtracting the gut fill from slaughter body weight. Hot carcass weight was computed by excluding contents of thoracic, abdominal and pelvic cavities, head, skin with feet (cut off at the proximal end of cannon bone) and tail of the animal. Dressing percentage was calculated as a ratio of hot carcass weight to slaughter weight and empty body weight multiplied by 100. The rib-eye muscle area of each animal was determined by tracing the cross sectional areas of the 12th and 13th ribs after cutting perpendicular to the back bone. The left and right rib-eye muscle areas were traced on a transparent water proof paper and the area was calculated as recommended by (Torell and Soverly, 2004). The value for rib-eye muscle area was the average of the right and left sides on 12th rib. Total edible offal (TEO) components were taken as the sum of blood, testicles, liver, kidney, kidney fat, heart, omental fat, tongue, reticulo-rumen, omasum-abomasum, large and small intestine and tail. Total non-edible offal components (TNEO) were computed as the sum of spleen, pancreas, skin and feet with hooves, penis, head without tongue, lung with trachea and gut contents. Total useable products (TUP) were taken as the sum of hot carcass weight, TEO and skin.

Statistical Data Analysis: The data generated during the experimental period were subjected to statistical analysis of variance (ANOVA) following the General Linear Model (GLM) procedure of SAS version 9.0 (SAS, 2002). Significant means ($P < 0.05$) were separated by Duncan's multiple range tests.

The model used for data analysis was:

$$Y_{ij} = \mu + t_i + b_j + e_{ij}$$

Where; Y_{ij} = response variable, μ = overall mean, t_i = treatment effect, b_j = block effect, e_{ij} = random error

RESULTS AND DISCUSSION

Carcass Characteristics

Main carcass components: There were significant effects in relation to increasing the levels of Dolichos lablab inclusion in the Elephant grass silage on slaughter weight (SW) ($P < 0.01$), empty body weight (EBW), hot carcass weight (HCW) ($P < 0.001$), Hind quarter, fore quarter, dressing percentage both on slaughter and empty body weight basis and rib eye muscle

area (REMA) ($P < 0.05$) all showed increased with increasing inclusion levels of Dolichos lablab in the Elephant grass silage. Different studies have showed that supplementation had significant effect on slaughter weight, empty body weight, hot carcass weight, rib-eye muscle area (Abebe, 2006; Matiwas *et al.*, 2008; Wondesen, 2008; Tewodros, 2011; Mulugeta and Gebrehiwot, 2013). Similarly, Mekonnen *et al.*, (2017) reported that Horro lambs fed *Cynodon dactylon* hay with nitrogen rich leguminous forage has significantly improved the parameters indicated. The present study result for dressing percentage on slaughter and empty body weight basis was comparable with the values noted by Assefu (2012) for Horro sheep weighing 21.2 to 24.7 Kg at slaughter ranged from 35 to 36.2% and 42.2 to 44%, by Mekonnen *et al.* (2017) for the same breed of 22.2 to 27.5 Kg slaughter weight ranged from 34.2 to 37.5% and 46 to 47.5% respectively.

Rib-eye muscle area is mostly used as a tool to indicate the proportion of carcass muscling (Wolf *et al.*, 1980). In the present study, Rib-eye muscle area was significantly ($P < 0.05$) influenced by dietary treatment. According to the report of Wolf *et al.* (1980), greater rib-eye muscle area is associated with a higher production of lean in the carcass and higher lean to bone ratio. Lambs assigned to T4 diet showed relatively higher REMA, which is a reflection of increase in lean meat. In the current study slightly higher rib-eye muscle area ranged from 6.4 to 9.93cm² was obtained as compared to 7.4 to 8.9 cm² reported by (Assefu, 2012). Comparable values for rib-eye muscle area 6.0 to 9.5cm² were reported by Mekonnen *et al.* (2017) for the same breed of sheep fed *Cynodon dactylon* hay alone or supplemented with *Cajanus cajan*, *Lablab purpureus* or their mixture. Similarly, Abuye *et al.* (2018b) noted rib eye area ranged from 7.04-9.54 cm² for the same breed of sheep fed natural grass hay and supplemented with Gebisa-17 and Beresa-55 cultivars of *Lablab purpureus* and concentrate mixture.

Non carcass parameters: Offal components are edible or non-edible based on tradition, religion, culture and differences in norms and preferences of the people and is more or less subjective.

Edible offal components: The mean weight of blood, kidneys, heart fat, reticulo-rumen and tongue of lambs fed Elephant grass ensiled with different proportion of Dolichos lablab were not affected Table 6. The weight mean values of tail, heart, liver with gall bladder, omasum-abomasum, omental fat, kidney fat, testicles, Large and small intestine, and total edible offal components (TEOC) were significantly influenced ($P < 0.05$) by dietary treatments. Similar to the present study, Mekonnen *et al.* (2017) reported that the weight of carcass offal such as heart, liver with gall bladder, intestines and tail is significantly improved by supplementation in Horro lambs fed *Cynodon dactylon* hay supplemented with *Cajanus cajan*, *Lablab purpureus* or their mixture. The values of large and small intestine, heart and tail weight was significantly higher ($P < 0.05$) for lambs fed T4 diet as compared to T1, but it is not significantly different ($P > 0.05$) from those fed diets in T2 and T₃. In the present study the value of liver with gall bladder weight was significantly higher ($P < 0.05$) for lambs fed diets in T4 as compared to those in T1 and T2 but had values similar ($P > 0.05$) with those in T3 group. Various studies Tewodros (2011); Mekonnen *et al.* (2017) and Abuye *et al.* (2018b) reported an increasing trend in weight of liver with

Table 1. Chemical composition of experimental feeds

| Feed Samples | DM% | Chemical Composition, (% DM) | | | | | | |
|-----------------|-------|------------------------------|-------|-------|-------|-------|------|--------|
| | | Ash | CP | OM | NDF | ADF | ADL | IVDOMD |
| T1 | 30.14 | 10.86 | 8.75 | 89.14 | 72.14 | 42.23 | 8.77 | 55.1 |
| T2 | 32.40 | 8.64 | 10.19 | 91.36 | 63.00 | 40.89 | 4.56 | 56.24 |
| T3 | 34.70 | 8.21 | 11.06 | 91.79 | 61.11 | 38.86 | 4.02 | 56.55 |
| T4 | 35.34 | 7.83 | 12.38 | 92.17 | 61.21 | 36.19 | 4.32 | 58.87 |
| Wheat Bran (WB) | 90.76 | 6.09 | 17.37 | 93.91 | 37.36 | 15.21 | 3.88 | 78.61 |
| Refusal | | | | | | | | |
| T1 | 30.08 | 13.1 | 6.48 | 86.9 | 70.86 | 42.51 | 9.32 | 36.65 |
| T2 | 30.83 | 12.58 | 6.87 | 87.42 | 72.82 | 41.68 | 4.63 | 38.19 |
| T3 | 31.44 | 12.52 | 7.13 | 87.48 | 71.64 | 40.86 | 4.59 | 39.81 |
| T4 | 31.92 | 11.23 | 7.25 | 88.77 | 71.6 | 40.79 | 4.59 | 40.43 |

DM= dry matter; OM= organic matter; NDF=neutral detergent fiber; ADF=acid detergent fiber; ADL= acid detergent lignin; CP= crude protein; IVDOMD= In vitro digestible organic matter in dry matter; NG= Elephant grass; DL= Dolichos lablab; WB = wheat bran; T1=Elephant grass ensiled alone; T2= Elephant grass ensiled with 10% Dolichos Lablab; T3=Elephant grass ensiled with 20% Dolichos Lablab; T4=Elephant grass ensiled with 30% Dolichos Lablab.

Table 2. Carcass components of Horro lambs fed Elephant grass ensiled with different proportion of Dolichos lablab

| Parameters | Treatments | | | | SEM | SL |
|---------------------------------|--------------------|---------------------|---------------------|--------------------|--------|-----|
| | T1 | T2 | T3 | T4 | | |
| Slaughter Weight (Kg) | 22.92 ^b | 24.96 ^{ab} | 25.96 ^a | 27.68 ^a | 0.6108 | ** |
| Empty Body Weight (Kg) | 18.58 ^c | 20.24 ^{bc} | 21.57 ^{ab} | 22.24 ^a | 0.3890 | *** |
| Hot Carcass Weight (Kg) | 7.78 ^b | 8.37 ^b | 9.96 ^a | 10.89 ^a | 0.3948 | *** |
| Fore Quarter (Kg) | 3.96 ^b | 4.64 ^{ab} | 5.22 ^{ab} | 5.83 ^a | 0.2887 | ** |
| Hind Quarter (Kg) | 3.48 ^b | 4.22 ^{ab} | 4.94 ^{ab} | 5.04 ^a | 0.3603 | * |
| Rib-eye area (cm ²) | 6.4 ^b | 7.53 ^b | 7.90 ^{ab} | 9.93 ^a | 0.7581 | * |
| Dressing Percentage on | | | | | | |
| Slaughter Weight basis | 33.94 ^b | 33.54 ^b | 38.36 ^a | 39.23 ^a | 1.509 | * |
| Empty Body Weight basis | 41.70 ^b | 41.41 ^b | 46.19 ^{ab} | 48.89 ^a | 1.618 | * |

^{a,b,c,d} =Means within a row not bearing a common superscript differ significantly; *=(P<0.05); **=(P<0.01); ***=(P<0.001); SEM=standard error of means; SL= significance level; T1 to T4= treatments..

Table 3. Edible offal of Horro lambs fed Elephant grass ensiled with different proportion of Dolichos lablab

| Edible Offal Components (g) | Treatments | | | | SEM | SL |
|-----------------------------|----------------------|-----------------------|----------------------|----------------------|--------|-----|
| | T1 | T2 | T3 | T4 | | |
| Blood | 1070 | 1040 | 1234 | 1150 | 62.007 | NS |
| Tail | 422 ^b | 654 ^{ab} | 694.4 ^{ab} | 806 ^a | 83.317 | * |
| Heart | 99.2 ^b | 104.46 ^{ab} | 117.4 ^{ab} | 134.8 ^a | 8.1304 | * |
| Heart fat | 50.12 | 43.32 | 52.58 | 50.2 | 3.4505 | NS |
| Liver with gall bladder | 388.38 ^c | 406.84 ^{bc} | 447.86 ^{ab} | 461.76 ^a | 14.887 | * |
| Kidneys | 127.8 | 117.86 | 119.28 | 129.36 | 8.8129 | NS |
| Kidneys fat | 37.72 ^c | 42.46 ^{bc} | 49.64 ^b | 59.40 ^a | 2.3676 | *** |
| Large and Small Intestine | 830 ^b | 992 ^{ab} | 1196 ^a | 1092 ^a | 74.030 | * |
| Reticulo-rumen | 592.2 | 682 | 604 | 568 | 36.270 | NS |
| Omasum-abomasum | 208 ^c | 360 ^a | 270 ^{bc} | 286 ^b | 21.726 | ** |
| Omental fat | 24.80 ^c | 27.70 ^{bc} | 30.06 ^b | 34.54 ^a | 1.2572 | *** |
| Testicles | 236.2 ^b | 221.98 ^a | 261.4 ^a | 329.6 ^a | 19.107 | ** |
| Tongue | 88.80 | 87.12 | 95.92 | 92.48 | 3.5334 | NS |
| Total Edible offal | 4175.22 ^b | 4779.74 ^{ab} | 5172.54 ^a | 5194.14 ^a | 171.16 | ** |

^{a,b,c,d} =Means within a row not bearing a common superscript differ significantly; *=(P<0.05); **=(P<0.01); ***=(P< 0.001); SEM=standard error of means; SL= significance level; T1 to T4= treatments.

Table 4. Non Edible offal of Horro lambs fed Elephant grass ensiled with different proportion of Dolichos lablab

| Non-Edible Offal Components (g) | Treatments | | | | SEM | SL |
|---------------------------------|---------------------|--------------------|--------------------|--------------------|--------|-----|
| | T1 | T2 | T3 | T4 | | |
| Skin | 1976 ^b | 2150 ^{ab} | 2280 ^a | 2414 ^a | 76.875 | * |
| Gut contents | 4985.2 ^a | 4482 ^a | 3912 ^b | 3770 ^b | 192.93 | ** |
| Head without Tongue | 1210 ^b | 1384 ^{ab} | 1420 ^a | 1550 ^a | 71.366 | * |
| Lung with trachea | 290.64 | 317.6 | 310.6 | 313.72 | 14.333 | NS |
| Spleen | 45.3 | 43.72 | 49.62 | 46.46 | 2.4067 | NS |
| Pancreas | 30.06 | 32 | 35.74 | 34.36 | 1.9877 | NS |
| Penis | 45.76 | 52.38 | 53.1 | 46.26 | 3.0244 | NS |
| Feet with hooves | 472 | 484 | 500 | 517.6 | 14.324 | NS |
| Total Non-Edible offal | 9055 | 8945 | 8561.1 | 8692.4 | 287.32 | NS |
| Total Usable Products (Kg) | 13.95 ^b | 15.25 ^b | 17.43 ^a | 18.49 ^a | 0.4537 | *** |

^{a,b,c,d} =Means within a row not bearing a common superscript differ significantly; *=(P<0.05); **=(P<0.01); ***=(P< 0.001); SEM=standard error of means; SL= significance level; T1 to T4= treatments.

increase in the supplementary diets and suggested that this phenomenon could happen due to the storage of more reserve substances such as glycogen in the liver. Gebregziabher *et al.* (2003) also reported higher liver weight in Horro sheep fed with sesbania forage compared to those fed ground maize grain and grass hay, which supports the present result. Similarly, the weight of omasum-abomasum, testicles and total edible offal component was significantly different ($P < 0.01$) among dietary treatments. Lower weight of testicles, omasum-abomasum and total edible offal for sheep fed T1 diet as compared to those fed T4, T3 and T2 diet. While, significantly higher ($P < 0.001$) values of omental fat and kidneys fat weight was observed for sheep fed T4 diet as compared to those in other group.

In general, increasing level of Dolichos lablab inclusion in Elephant grass silage affected positively the omental fat and other edible offal components production. This might be due to the high energy content of legume enriched silage feed which promoted higher internal fat deposition. Gameda *et al.* (2002) reported that testis weight differences between the animals in the different treatments may reflect the body weight differences among the different treatments. Significantly increased TEO with increasing level of supplementation for Horro sheep fed natural grass hay and supplemented with Gebisa-17 and Beresa-55 cultivars of *Lablab purpureus* and concentrate mixture was reported by (Abuye *et al.*, 2018b). The author further reported that, TEO values ranging from 3.75 to 4.89 Kg which agrees with value obtained in the current study (4.175 to 5.194 Kg). Similarly, Comparable TEO values ranged from 3.08 to 4.9 Kg for the same breed was reported by (Girma and Mengistu, 2017). However, Assefu (2012) was reported the values of TEO weight ranged from 3.68 to 4.55 Kg which was slightly lower than the present values of TEO weight. Higher weight of TEO ranged between 4.3 and 5.8 Kg was reported by (Mekonnen *et al.*, 2017) for the same breed.

The values of Skin, gut contents and head without tongue were significantly influenced by feeding of ensiled Elephant grass with different proportion of Dolichos lablab. Whereas, the rest parameters indicated in (Table 7) were not influenced. In the present study, lambs assigned to diets in T1 had lighter ($P < 0.05$) skin and head without tongue as compared to lambs in other treatment group. Whereas, lambs fed diet in T2, T3 and T4 had almost similar ($P > 0.05$) value in their skin and head without tongue weight even though, lambs fed diet in T2 was not significantly different ($P > 0.05$) from T1. The relatively higher skin weight obtained by lambs fed diets in T4 and T3 might be associated to their higher ME intake of T4 (9.99 MJ/day) and T3 (9.41 MJ/day), which might have resulted in better subcutaneous layer fat deposition in these groups. The highest value of gut contents weight were recorded for lambs assigned to fed T1 and T2 diet. The heaviest gut content of lambs fed T1 and T2 diet may be due to the higher roughage or relatively poor quality feed used. This was agreed with the views of VanSoest (1994) and pond *et al.* (1995) in that non supplement animals fill their gut with less digestible roughage, which would retain in the gut for longer time to be degraded by rumen microbes. Total usable products showed highly significant ($P < 0.001$) variations among dietary treatments. It was significantly higher for lambs fed T4 and T3 diet as compared to those fed T1 and T2 diets.

Conclusion

The experiment was conducted at Bako Agricultural Research Center using intact male yearling Horro lambs with initial body weight of 19.47 ± 1.76 kg. The objectives of the study was to evaluate the effects of Elephant grass ensiled with different proportion of Dolichos lablab on carcass and non-carcass yield of Horro sheep. A randomized complete block design (RCBD) with four treatments and five replications were used. Animals in a blocks were assigned to one of the four treatments. Dietary treatments were arranged as T1, T2, T3 and T4 respectively for Elephant grass ensiled alone, with 10, 20 and 30% of Dolichos lablab. The experimental animals in all treatments were free access to clean water, mineralized salt block and supplemented with wheat bran of 1.5% body weight. All experimental animals were slaughtered for carcass characteristics evaluation.

Variations among experimental diets were observed in chemical composition. Dry matter and crude protein content of the silage increased as the proportion of Dolichos lablab in the mixture increased. On the contrary, Ash, NDF, ADF and ADL of the ensiled mixtures reduced showing that addition of Dolichos lablab was effective in improving nutrient values of the Elephant grass silage. The mean weight of blood, kidneys, heart fat, reticulo-rumen and tongue of lambs fed Elephant grass ensiled with different proportion of Dolichos lablab were not affected. The weight mean values of tail, heart, liver with gall bladder, omasum-abomasum, Large and small intestine, head with tongue and total edible offal components (TEOC) were significantly influenced ($P < 0.05$) by dietary treatments. The weight values of Skin, gut contents and head without tongue of Horro lambs were significantly influenced by feeding of ensiled Elephant grass with different proportion of Dolichos lablab whereas, the rest parameters including lung with trachea, spleen, pancreas, penis and feet with hooves were not influenced. In general, based on the result, it can be concluded that mixing Dolichos lablab with Elephant grass had beneficial effects on silage properties and carcass yield of Horro lamb. Therefore, it is recommended that addition of 20-30% Dolichos lablab enhances the nutritive value of Elephant grass silage thereby, increasing the growth and carcass yield of Horro sheep.

REFERENCES

- Abebe Tefa. 2006. Supplementation with linseed (*linum usitatissimum*) cake, wheat bran and their mixtures on feed intake, digestibility, live weight changes, and carcass characteristics in intact male Arsi-Bale sheep. MSc. Thesis Presented to School of graduate studies of Haramaya University, Haramaya, Ethiopia.65p.
- Abuye Tullu, Yadav, R.K. and Diriba Geleti. 2018a. Carcass and non-carcass yield characteristics of horro sheep supplemented with two lablab purpureus cultivars and concentrate mixture to a basal diet of natural grass hay. *Journal of Biology, Agriculture and Healthcare*. 8(13): 2224-3208.
- Abuye Tullu, Yadav, R.K. and Diriba Geleti. 2018b. Supplementary value of two *Lablab purpureus* cultivars and concentrate mixture to natural grass hay basal diet based on feed intake, digestibility, growth performance and net return of Horro sheep. *International Journal of Livestock Production*. 9(6): 140-150.

- Adegun, M.K. 2014. Voluntary Feed Intake and Nutrient Utilization of West African Dwarf Sheep Fed supplements of *Moringa oleifera* and *Gliricidia sepium* Fodders. *American Journal of Agriculture and Forestry*, 2(3): 94-99.
- Adesogan, A.T., Salawu, M.B., Williams, S.P., Fisher, W.J. and Dewhurst, R.J. 2002. Reducing concentrate supplementation in dairy cow diets while maintaining milk production with pea-wheat intercrops. *Journal of Dairy Science*, 87: 3398–3406.
- Ajebu Nurfeta, Adugna Tolera, Eik, L.O. and Sundstol, F. 2008. Feeding value of ensiled (*Ensete ventricosum*), *Desmodium intortum* hay and untreated or urea and calcium oxide treated wheat straw for sheep. *Journal of Animal Physiology and Animal Nutrition*, 93: 94-104.
- Alemayehu Mengistu. 2002. Recommended Forage for Ethiopian Environments and Current Strategies for Utilization. Proceedings of the XVI International Grassland Congress, Nice, France.
- ARC (Agricultural Research Council). 1980. The Nutrient Requirements of Ruminant Livestock Technical review by an Agricultural Research Council Working party published on behalf of the Agricultural Research Council by the Common Wealth Agricultural Bureaux, Farnham Royal, England pp.114-151.
- Assefu Gizachew. 2012. Comparative feedlot performance of Washera and Horro sheep fed different roughage to concentrate ratio. An MSc Thesis Presented to School of graduate studies of Haramaya University, Haramaya, Ethiopia.68p.
- Awet Estifanos and Solomon Meleku. 2009. Supplementation of Graded Levels of Wheat Bran to Intact and Castrated Afar Sheep Fed Urea Treated Teff Straw: Effects on Feed Intake, Digestibility, Body Weight and Carcass Characteristics. *East African Journal of Science*, 3(1): 29-36.
- Banamana, M.S., Oldman, J.D. and Mowlem, A. 1990. The effect of amount of protein in the Concentrate on hay intake and rate of passages diet digestibility and milk production in British Saanen goat. *Journal of Animal Production*, 51:333-342.
- Belete Shenkute, Abubeker Hassen, Abule Ebro and Nura Amen. 2013. Performance of Arsi-Bale kids supplemented with graded levels of pigeon pea in dry season in Mid Rift valley of Ethiopia. *African Journal of Agricultural Research*, 8(20): 2366-2370.
- Berhanu Alemu, Getachew Animut and Adugna Tolera. 2014. Effect of *Milletia ferruginea* (Birbra) foliage supplementation on feed intake, digestibility, body weight change and carcass characteristics of Washera sheep fed natural pasture grass hay basal diet. *SpringerPlus*. pp.4-6.
- Birhanu T, Getachew Animut, Mengistu Urgei. 2013. Effect of Green *Prosopis juliflora* Pods and Noug Seed (*Guizotia obissynica*) Cake Supplementation on Digestibility and Performance of Blackhead Ogaden Sheep fed hay as a Basal Diet Science Technology Arts Research Journal, 2(2): 38-47.
- Bonsi, M.L.K., Tush, A.K., Osuji, P.O., Nsahlai, I.V. and Umunna, N.N. 1996. The effect of protein supplement source or supply pattern on the intake, digestibility, rumen kinetics, nitrogen utilization and growth of Ethiopian Menz sheep fed teff straw. *Journal of Animal Feed science and Technology*, 64(1): 11-25.
- Brown, L., Hindmarsh, R., McGregor, R. 2001. Dynamic Agricultural book three (2nd Ed.): McGraw-Hill Book Company, Sydney, 357p.
- Contreras-Govea, F.E., Muck, R.E., Armstrong, K.L. and Albrecht, K.A. 2009. Nutritive value of corn silage in mixture with climbing beans. *Animal Feed Science Technology*, 150: 1-8
- CSA. 2018 (Central Statistical Agency Agricultural) sample survey 2017/18. Report on Livestock and livestock characteristics (private peasant holdings) volume, II.
- DAGRIS (Domestic Animals generic Resources Information System). 2006. (Rege, J.E.O., Workineh Ayalew, Getahun, E., Hanotte, O. and Tadele Dessie) International livestock Research Institute, Addis Ababa Ethiopia [http:// dagris ilri.Cigar. Org.](http://dagris.ilri.org)
- Ensminger, M.E. 2002. Sheep and goat science (Animal Agriculture Series). 6th ed. Interstate publishers, Inc. Daville, Illinois, U.S.A. 693p
- Ferreira, D.J., Zanine, A.M., Lana, R.P., Souza, A.L., Ribeiro, M.D., Negrão, F.M., Alves, G.R. and Castro, W.J.R. 2017. Intake and digestibility in sheep fed on marandu grass silages added with dehydrated barley. *Pesquisa Veterinária Brasileira* 37(2):171-178.
- Firaol Edao. 2017. Effects of replacing different proportion of cafeteria leftover for concentrate mix on performance of Arsi – Bale sheep. A Thesis submitted to School of Animal and Range Sciences (Postgraduate Program Directorate) Haramaya University.
- Firisa, W., Adugna Tolera, Diriba Dida. 2013. Feed Intake, Digestibility and Growth Performance of Horro lambs fed Natural Pasture hay Supplemented Graded Level of *Vernonia amygdalina* Leaves and Sorghum Grain Mixture. *Science and Technology Arts Research Journal*, 2(2):30-37.
- Gebregziabher Gebreyohannis, Diriba Geleti, Lemma Gizachew, Yohannes Gojjem and Gemeda Duguma. 2003. Effect of noug cake and Sesbania Sesban supplementation on the growth performance and carcass characteristics of Horro rams, 335-339. Proceedings of the 10th annual conference of the Ethiopian Society of Animal Production (EASP), 21-23 August, Addis Ababa, Ethiopia.
- Gemeda Duguma, Cloete, S.W.P., Schoeman, S.J. and Jordaan, G.F. 2002. Genetic parameter estimates of testicular measurements in Merino rams and the influence of scrotal circumference on total flock fertility. *South Africa Journal of Animal Science*, 32:76–82.
- Gemeda Duguma, Takele Kumsa, Ulfina Galmessa and Solomon Abegaz. 2003. The effect of age and sex on growth performance and carcass characteristics of Horro lamb, 99-103. Proceedings of 10th Annual conference of the Ethiopian Society of Animal Production (ESAP). 22-24 August 2002, Addis Ababa, Ethiopia.
- Getnet Berhanu. 1998. Performance of Somali Goats Supplemented with Different Proportions of Groundnut Cake and Wheat bran. MSc Thesis, Haramaya University, Haramaya, Ethiopia.
- Girma Defar and Mengistu Urgie. 2017. Bodyweight gain and carcass characteristics of Horro sheep fed urea treated maize husk and untreated maize husk supplemented with different levels of concentrate mix at Bako, Western Ethiopia. *International Journal of Livestock Production*, 8(6): 87-94.
- Gulten, K., Rad, F. and Kindir, M. 2000. Growth performance and feed conversion efficiency of Siberian Sturges

- juveniles (*Acipenser baeri*) Reared in concentrate ways. *Turkey Journal of Veterinary and Animal Science*, 24:28.
- Hindrichsen, I.K., Osuji, P.O., Odenyo, A.A., Madsen, J. and Hvelplund, T., 2002. Effects of supplementation of a basal diet of maize stover with different amounts of *Leucaena diversifolia* on intake, digestibility, nitrogen metabolism and rumen parameters in sheep. *Animal Feed Science and Technology Journal*, 98:131-142.
- Howes, N.L., Bekhit, A.E.A., Burritt, D.J. and Campbell, A.W. 2015. Opportunities and implications of pasture-based lamb fattening to enhance the long-chain fatty acid composition in meat. *Comprehensive Review in Food Science and Food Safety*, 14:22–36.
- Hunagnaw Abebe. 2015. Effects of supplementation with pigeon pea (*cajanus cajan*), cowpea (*vigna unguiculata*) and lablab (*lablab purpureus*) on feed intake, body weight gain and carcass characteristics of wollo sheep fed on grass hay. MSc Thesis, Addis Ababa University, Addis Ababa. Ethiopia.
- Jalel, F. 2013. Effect of feeding graded levels of wild silver leaf desmodium (*Desmodium uncinatum*) on feed intake, digestibility, body weight change and carcass characteristics of Horro sheep fed a basal diet of Natural pasture hay. MSc Thesis, Haramaya University, Haramaya, Ethiopia, pp. 24-36.
- Jemal Dekebo, Chala Merera, Diriba Geleti, Ulfina Gelmessu, Diriba Diba and Adane Hirpa. 2004. Weight gain, carcass characteristics and economic evaluation of Horro rams supplemented with different substitution levels of maize grain and molasses. 45-51. Proceeding of the 13th annual conference of the Ethiopian Society of Animal Production (ESAP), 25-27 August 2004, Addis Ababa, Ethiopia.
- J rgensen, S.T., Pookpakdi, A., Tudsri S., Stölen O., Ortiz R. and Christiansen J.L. 2010. Cultivar-by-cutting height interactions in Elephant grass (*Pennisetum purpureum* Schumach) grown in a tropical rain-fed environment *Acta Agriculturae Scandinavica, Section B- Plant Soil Science* 60: 199-210.
- Keady, T.W.J. and O'Kiely, P. 1998. An evaluation of potassium and nitrogen fertilization of
- Lapenga, K. O., Ebong, C. and Opuda-Asibo, J. 2009. Effect of feed supplementation on weight gain and carcass characteristics of intact male Mubende goats fed Elephant grass (*Pennisetum purpureum*) ad libitum in Uganda. *Journal of animal and veterinary advances*, 8 (10): 2004-2008.
- Malede Birhan and Takele Adugna. 2014. Livestock feed resources assessment, constraints and improvement strategies in Ethiopia. *Middle East Journal of Scientific Research*, 21(4): 618–704.
- Malisetty, V., Yerradoddi, R.R., Devanaboina, N., Mallam, M., Cherala, H.K., Admal, R.R. and Manthani, G.P. 2013. Effect of feeding maize silage supplemented with concentrate and legume hay on growth in Nellore ram lambs, *World*. 6(4): 209-213, doi:10.5455/vetworld.2013.209-213
- Mapiye, C., Mwale, M., Mupangwa, J.F., Mugabe, P.H. Poshiwa, X. and Chikumba, N. 2007. Utilisation of ley legumes as livestock feed in Zimbabwe. *Tropical Grass*, 41: 84-91
- Markos Tibbo, Phillips, J.S. and Worknesh Ayalew. 2006. Sustainable sheep breeding in the Tropics: a Framework for Ethiopia conference on International Agricultural Research for Development, University of Bonn, October, 2006.
- Matiwos Solomon, Solomon Melaku and Adugna Tolera. 2008. The effect of different levels of cottonseed meal supplementation on feed intake, digestibility, live weight changes, and carcass parameters of Sidama Goats. *Livestock Science*, 119: 137-144.
- Gachuri, C.K. and Mbuthia, E.W. 2003. Effect of inclusion of *mucuna pruriens* and *Dolichos lablab* forage in Elephant grass silage on silage quality and on voluntary intake and digestibility in sheep. *Tropical and Subtropical Agroecosystems*, 1: 123–128.
- McDonald, P., Edwards, A.R., Greenhalgh, D.F.J. and Morgan, A.C. 2002. *Animal Nutrition*. 6th ed. Prentice Hall, London. 245-477.
- McDonald, P., Henderson, A.R. and Heron, S.J.E. 1991. The biochemistry of silage, 2nd Edition. Marlow, UK: Chalcombe Publications.
- Mekonnen Diribsa, Mengistu Urgie and Gemedu Duguma. 2016. Effects of supplementation with *cajanus cajan*, *lablab purpureus* or their mixture on feed utilization, growth and carcass characteristics of Horro sheep fed a basal diet of natural grass hay. *Journal of Biology, Agriculture and Healthcare*. 6(17): 2224-3208.
- Mekonnen Diribsa. 2017. Effect of supplementing improved forage legumes (*Cajanus cajan* and *Lablab purpureus*) on carcass characteristics of Horro sheep fed a basal diet of *cynodon dactylon* hay. *Journal of Natural Sciences Research*. 7(19): 2224-3186.
- Mhere, O., Maasdorp, B. and Titterton, M. 2002. Growing and ensiling annual and perennial forage crops suited to marginal and semi-arid areas of Southern Africa. *Forage Production and Conservation Manual*.
- Muia, J.M.K., Tamminga, S., Mbuguam, P.N. and Kariuki, J.N. 2001. Rumen degradation and estimation of microbial protein yield and intestinal digestion of Elephant grass (*Pennisetum purpureum*) and various concentrates. *Animal Feed Science and Technology*, 93: 177-192.
- Mulugeta, F. and Gebrehiwot, T. 2013. Effect of sesame cake supplementation on feed intake, body weight gain, feed conversion efficiency & carcass parameters in the ration of sheep fed on wheat bran & teff (*Eragrostis teff*) straw. *Momona Ethiopian Journal of Science*, 5(1): 89-106.
- Napasirth, V., Napasirth, P., Sulinthone, T., Phommachanh, K. and Cai, Y. 2015. Microbial population, chemical composition and silage fermentation of cassava residues. *Animal Science Journal*, 86: 842–848.
- Ndikumana, J. 1996. Multi locational evaluation of Elephant and *Pennisetum* hybrids: Highlights of from the Addis Ababa. AFRNET (Africa Feed Resources Network) newsletter, 6: 1-11.
- Ngongoni, N.T., Mwale, M., Mapiye, C., Moyo, M.T., Hamudikuwanda, H. and Titterton, M. 2008. Evaluation of cereal-legume intercropped forages for smallholder dairy production in Zimbabwe. *Tropical Grassland Science*, 42: 188–192.
- Olorunnisomo, O.A. 2011. Intake and digestibility of elephant grass ensiled with cassava peels by Red Sokoto goats. Conference on International Research on Food Security, Natural Resource Management and Rural Development. University of Bonn, October 5-7, 2011.
- Osuji, P.O., Fernandez-Rivera, S. and Odenyo, A.A. 1995. Improving fiber utilization and protein supply in animals fed poor quality roughages: ILRI nutrition research and

- plans. 1-22. Rumen Ecology Research Planning. Addis Ababa, Ethiopia, In: 13-18 March 1995, ILRI.
- Pahlow, G., Slottner, D., Rammer, C. and Touri, M. 2002. Ensiling of legumes. Published in: Landbauforschung Völkenrode Sonderheft 234, PP. 27-31.
- Phelan, P., Moloney, A.P., McGeough, E.J., Humphreys, J., Bertilsson, J., O'Riordan, E.G. and O'Kiely, P. 2015. Forage legumes for grazing and conserving in ruminant production systems. *Critical Reviews in Plant Sciences*, 34(13): 281–326.
- Radia Hussien, Firew Tegegne, Zelalem Yilma, Zeleke Mekuriaw, Mengistie Taye. 2013. Feed Intake, Milk Yield and Milk Composition of Fogera Cows Supplemented with Different Feeds. *World's Veterinary Journal*, 3(2): 41 - 45.
- Ranjihan, S.K. 2001. Animal nutrition in the Tropics. 5th revised edition Vikas publishing Resources Information. 43: 25–40.
- SAS (Statistical Analysis System). 2002. User's Guide: version 9.0. SAS Institute, Inc. Cary, NC.
- Seyoum Bediye and Zinash Sileshi, 1998. The composition of Ethiopian feed stuffs, Research report No.6. Institute of agriculture research (IAR), Addis Ababa, Ethiopia. 33p.
- Simret Betsha. 2005. Supplementation of graded levels of peanut cake and Wheat bran mixture on nutrient utilization and Carcass parameters of Somali goats. Thesis presented to the School of Graduate Studies of Haramaya University. Haramaya, Ethiopia. 75p
- Solomon Gizaw, Solomon Abegaz, Rischkowsky B. Haile, A., Mwai, A.O. and Tadele Dessie. 2013. Review of sheep research and development projects in Ethiopia. Nairobi, Kenya: International Livestock Research Institute (ILRI).
- Solomon Melaku, Peters, K.J. and Azage Tegegne. 2004. Effects of supplementation with foilages of selected multipurpose trees, their mixtures or wheat bran on feed intake, plasma enzyme activities, live weight and scrotal circumference gain in Menz sheep. *Livestock Production Science*, 89: 253-264.
- Susan, S. 2003. An introduction to feeding small ruminants. Western Maryland Research and Education Center Maryland Cooperative Extension. <http://www.sheepandgoat.com/index.html>
- Takele Kumsa, Gameda Duguma, Fayo Dubiso and Solomon Abegaz. 2004. Growth performance of Horro lambs supplemented with different levels of wilted *Leucaena (Leucaena Pallida)* leaf. pp.248-252. Proceedings of the 12th annual conference of the Ethiopian Society of Animal Production (ESAP), 12-14 August 2004, Addis Ababa, Ethiopia.
- Teklu Wegi. 2016. Effects of feeding different varieties of faba bean (*vicia faba* l.) straws with concentrate on feed intake, digestibility, body weight gain and carcass characteristics of Arsi-Bale sheep. MSc Thesis, Haramaya University, Haramaya, Ethiopia.
- Tesfaye Hagos. 2007. Supplementation of Afar rams with graded levels of mixtures of protein and energy sources: Effects on feed intake, digestibility, live weight and carcass parameters. An MSc Thesis presented to School of Graduated Studies Haramaya University, Haramaya, Ethiopia. 24p.
- Tessema Zewdu. 2005. Variation in growth, yield, chemical composition and *in vitro* dry matter digestibility of Elephant grass accessions (*Pennisetum purpureum*). *Tropical Science*, 45: 93-99.
- Tessema, Zewdu, Ashagre, A. and Solomon Melaku. 2010. Botanical composition, yield and nutritional quality of grassland in relation to stages of harvesting and fertilizer application in the highland of Ethiopia. *African Journal of Range and Forage Science*.
- Tewodros Eshete. 2011. Effect of inclusion of different proportions of tosign (*thymus serrulatus*) in concentrate mix supplement on feed intake, digestibility, body weight change and carcass parameters of Menz sheep fed grass hay. MSc Thesis Presented to the School of Graduate Studies of Haramaya University, Haramaya, Ethiopia. 57p.
- Titterton, M. and Bareeba, F. 1999. Paper 4.0: Grass and legume silages in the tropics. FAO, Rome, Retrieved from <http://www.fao.org/docrep/005/x8486e/x8486e0c.htm> [Cited 23 October 2016].
- Torell, R. and Suverly, N. 2004. Nevada market lamb carcass merit program. Fact sheet 04 02.
- Upton, M. 1979. Farm management in Africa: the principle of production and planning. Oxford University press, Great Britain. pp. 282-298.
- Van Soest, P.J., Robertson, B.J. and Lewis, B.A. 1994. Methods for dietary fiber, neutral detergent fiber, and non-starch polysaccharides in relation to animal nutrition. *Journal of Dairy Science*, 74: 3583-3597.
- Wakene Negassa. 2001. Assessment of important physicochemical properties of Alfisols under different management systems in Bako area, Western Ethiopia. An MSc Thesis Presented to the School of Graduate Studies of Alemeya University, Haramaya, Ethiopia. 93p.
- Wolf, B.T., Smith, C. and Sales, D.I. 1980. Growth and carcass composition in the crossbred Progeny of six terminals sire breeds of sheep. *Animal Production*, 31: 307-313.
- Wondwesen Bekele. 2010. Effect of substitution of concentrate mix with *Sesbania sesban* on feed intake, digestibility, body weight change, and carcass parameters of Arsi-Bale sheep fed grass hay. MSC Thesis, Haramaya University, Haramaya, Ethiopia.
- Wondwosen Alemu. 2008. Effect of supplementing hay from natural pasture with oil seed cakes on feed intake, digestibility and live weight change of Sidama goats. An MSc. Thesis Presented to School of Graduate Studies of Haramaya University. 27p.
- Wong, C.C. 2000. The place of silage in ruminant production in the humid tropics. In: Silage Making in the Tropics with Particular Emphasis on Smallholders ('t Mannetje L., Ed.)
- Yenesew Abebe, Solomon Melaku and Azage Tegegne. 2013. Effect of supplementation of grazing sheep with groundnut cake and wheat bran. *Journal of Agricultural Research*, 2(8): 212-217.
- Yeshambel Mekuriaw, Mengistu Urge and Getachew Animut. 2012. Intake, digestibility, live weight changes and rumen parameters of Washera sheep fed mixtures of lowland bamboo (*oxytenanthera abyssinica*) leaves and natural pasture grass hay at different ratios. *Pakistan Journal of Nutrition*, 11 (4): 322-331.