



ISSN: 0976-3376

Available Online at <http://www.journalajst.com>

ASIAN JOURNAL OF  
SCIENCE AND TECHNOLOGY

Asian Journal of Science and Technology  
Vol. 12, Issue, 11, pp.11956-11959, November, 2021

## RESEARCH ARTICLE

### SEASONAL METABOLIC VARIATION OF PROTEIN and GLYCOGEN IN FRESHWATER BIVALVE LAMELIDENS CORRIANUS (LEA, 1834)

\*Dr. Suvarnamala K. Padewar

C.H.B. Lecturer, P.G. Department of Zoology, NES Science College, Nanded

#### ARTICLE INFO

##### Article History:

Received 14<sup>th</sup> August, 2021  
Received in revised form  
18<sup>th</sup> September, 2021  
Accepted 07<sup>th</sup> October, 2021  
Published online 28<sup>th</sup> November, 2021

##### Key words:

Protein, Glycogen,  
Lamellidens, Corrianus, Seasonal.

#### ABSTRACT

Proteins are building blocks. It is important organic substances required by an organism in tissues building and repair. Under extreme stress conditions, proteins have been known to act as the energy supplier in metabolic pathways and biochemical reactions. In the present investigation Protein is found maximum in gonads throughout summer season, increasing in rainy and minimum in winter season followed by mantle, hepatopancreas, Add. Muscle and gill i.e., Gonad> Mantle> Hepatopancreas> Add. muscle> Gill. Glycogen are important organic substances required by an organism in cellular process. Under extreme stress conditions, glycogen have been known to act as the energy supplier in metabolic pathways and biochemical reactions. In the present investigation glycogen is found maximum in gonads throughout summer season, increasing in rainy and minimum in winter season followed by mantle, hepatopancreas, Add. Muscle and gill i.e., Gonad> Mantle> Hepatopancreas> Add. muscle>Gill.

Citation: Dr. Suvarnamala K. Padewar, 2021. "Seasonal metabolic variation of protein and glycogen in freshwater bivalve lamellidens corrianus (lea, 1834)", *Asian Journal of Science and Technology*, 12, (11), 11956-11959.

Copyright © 2021, Suvarnamala K. Padewar. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Proteins are most abundant intracellular macro-molecules and constitute over half the dry weight of most organisms. They occupy a central position in the architecture and functioning of living matter. They are intimately connected with all phases of chemical and Physical activity that constitutes the life of the cell. Therefore they are, essential to cell structure and cell function. The interplay between enzymatic and non-enzymatic proteins governs the metabolic harmony (Lehinger, 1984). They are also involved in major physiological events to maintain the homeostasis of the cell. Therefore, the assessment of the protein content can be considered as a diagnostic tool to determine the physiological process of the cell (Kapil and Ragothaman, 1999; Munshigeri, 2003). Proteins are essential nutrients for the human body. They are one of the building blocks of body tissue, and can also serve as a fuel source. As a fuel, proteins contain 4 kcal per gram, just like carbohydrates and unlike lipids, which contain 9 kcal per gram. Proteins are polymer chains made of amino acids linked together by peptide bonds. During human digestion, proteins are broken down in the stomach to smaller polypeptide chains via hydrochloric acid and protease actions.

This is crucial for the synthesis of the essential amino acids that cannot be biosynthesized by the body (Genton *et.al.*, 2010). The chief carbohydrate in the tissues is glycogen, while glucose is an utilizable sugar found in the tissues and the body fluids. Glycogen is reversibly converted to glucose under the influence of hormonal mediated enzyme activities. The equilibrium in glycogenesis and glycogenolysis tends to maintain blood sugar at a steady state. The oxidation of glucose is mediated by catabolic pathways viz., glycolysis, Krebs cycle, electron transport system and hexose mono-phosphate shunt, which constitute the major segments of carbohydrate metabolism. Thus carbohydrate metabolism gained importance in the physiology of animals. According to Prosser, (1984) the synthesis and degradation of glycogen will not occur simultaneously at any significant rate. The other probability for occurrence of depletion in glycogen levels might be due to dephosphorylation of phosphorylase 'a' and specific protein phosphatases. Glycogen is a multibranched polysaccharide of glucose that serves as a form of energy storage in animals (Sadava *et. al.*, 2011). The polysaccharide structure represents the main storage form of glucose in the body.

## MATERIAL AND METHOD

Bivalves *Lamellidens corrians* sample (75-80 mm in shell length) were obtained from fishermen's catch. In the present study they were collected from Godavari River (Nanded) of

\*Corresponding author: Dr. Suvarnamala K. Padewar,  
C.H.B. Lecturer, P.G. Department of Zoology, NES Science College,  
Nanded.




Fig. Monthly variation in protein content of *Lamellidens corrianus*




Fig. Monthly variation in glycogen content of *Lamellidens corrianus*

Maharashtra in India in the year 2014. Immediately after bringing to laboratory, hard shells of these freshwater bivalves were brushed and washed with fresh and clean water to remove algal biomass, mud and other waste material. The cleaned animals were then kept for depuration for 12 hrs in laboratory conditions under constant aeration.

For protein analysis, animals were dissected and soft body tissues like mantle, hepatopancreas, gonad, Adductor muscle and gill were removed. 100 mg of each wet tissues were taken for biochemical analysis. Protein was determined by the method proposed by Lowry et al., (1951), using Bovine Serum Albumin (BSA) as standard and values of proteins were expressed in terms of mg protein/gm wet weight of tissue. Same procedure we have to do upto i.e. 100 mg of each wet tissues were taken for biochemical analysis. Then estimation of glycogen was done by Anthrone method using glucose as standard (Seifer *et al.*, 1950) and values of glycogen were expressed in terms of mg glycogen/gm wet weight of tissue.

## RESULTS AND DISCUSSION

Protein estimation observed during experimental work has been given in Fig for *Lamellidens corrianus*.

### Protein content in *Lamellidens corrianus*:

- **Gonad protein:** The seasonal changes in the protein content in gonad of *L. corrianus* are shown in (Table No. 4 and Figure No. 4). The percentage of protein was found to be maximum in summer and varies from  $10.1210 \pm 0.6301$  to  $12.0238 \pm 0.3332$ , increasing in rainy season and varies from  $6.1203 \pm 0.3412$  to  $10.1320 \pm 0.2103$ , whereas it is minimum in winter season and varies from  $4.2017 \pm 0.5204$  to  $6.1201 \pm 0.2360$ .
- **Mantle protein:** - The seasonal changes in the protein content in mantle of *L. corrianus* are shown in (Table No. 4 and Figure No. 4). The percentage of protein was found to be maximum in summer and varies from  $10.1232 \pm 0.3203$  to  $12.1022 \pm 0.4100$ , increasing in rainy season and varies from  $4.1310 \pm 0.4210$  to  $8.2162 \pm 0.5103$ , whereas it is minimum in winter season and varies from  $3.1231 \pm 0.1324$  to  $5.1130 \pm 0.3335$ .
- **Hepatopancreas protein:** The seasonal changes in the protein content in hepatopancreas of *L. corrianus* are shown in (Table No. 4 and Figure No. 4). The percentage of protein was found to be maximum in summer and varies  $7.1410 \pm 0.4250$  to  $10.1323 \pm 0.3504$ , increasing in rainy season and varies from  $5.1422 \pm 0.5140$  to  $7.3413 \pm 0.3312$ , whereas it is minimum in winter season and varies from  $3.2112 \pm 0.2310$  to  $6.2340 \pm 0.3221$ .
- **Add. Muscle protein:** The seasonal change in the protein content in Add. Muscle of *L. corrianus* are shown in (Table No. 4 and Figure No. 4). The percentage of protein was found to be maximum in summer and varies from  $9.3103 \pm 0.3230$  to  $11.1010 \pm 0.2310$ , increasing in rainy season and varies from  $8.1101 \pm 0.1012$  to  $7.1230 \pm 0.2031$ , whereas it is minimum in winter season and varies from  $3.4013 \pm 0.4530$  to  $5.1302 \pm 0.5230$ .
- **Gill protein:** The seasonal changes in the protein content in gill of *L. corrianus* are shown in (Table No. 4 and Figure No. 4). The percentage of protein was found to be maximum in summer and varies from  $8.1231 \pm 0.3323$  to  $9.4522 \pm 0.2520$ , increasing in rainy season and varies from  $7.6320 \pm 0.5360$  to  $8.5235 \pm 0.6320$ , whereas it is minimum in winter season and varies from  $2.3131 \pm 0.2221$  to  $5.3233 \pm 0.4014$ .

### Glycogen content in *Lamellidens corrianus*

- **Gonad glycogen:** - The seasonal changes in the glycogen content in gonad of *L. corrianus* are shown in (Table No. 5 and Figure No. 5). The percentage of glycogen was found to be maximum in summer and varies from  $10.0463 \pm 0.2352$  to  $11.1121 \pm 0.2342$ , increasing in rainy season and varies from  $4.3302 \pm 0.5203$  to  $9.1012 \pm 0.3414$ , whereas it is minimum in winter season and varies from  $3.1317 \pm 0.3204$  to  $5.0400 \pm 0.2231$ .
- **Mantle glycogen:** - The seasonal changes in the glycogen content in mantle of *L. corrianus* are shown in (Table No. 5 and Figure No. 5). The percentage of glycogen was found to be maximum in summer and varies from  $10.1213 \pm 0.7415$  to  $12.1220 \pm 0.1202$ , increasing in rainy season and varies from  $2.2210 \pm 0.5213$

- to  $6.6413 \pm 0.4201$ , whereas it is minimum in winter season and varies from  $6.6413 \pm 0.4201$  to  $6.6413 \pm 0.4201$ .
- **Hepatopancreas glycogen:** The seasonal changes in the glycogen content in hepatopancreas of *L. corrianus* are shown in (Table No. 5 and Figure No. 5). The percentage of glycogen was found to be maximum in summer and varies from  $10.1000 \pm 0.2110$  to  $11.0120 \pm 0.3220$ , increasing in rainy season and varies from  $4.1121 \pm 0.4120$  to  $7.4041 \pm 0.4403$ , whereas it is minimum in winter season and varies from  $4.2021 \pm 0.3210$  to  $6.2134 \pm 0.3010$ .
  - **Add. Muscle glycogen:** The seasonal changes in the glycogen content in Add. Muscles of *L. corrianus* are shown in (Table No. 5 and Figure No. 5). The percentage of glycogen was found to be maximum in summer and varies from  $9.0220 \pm 0.3130$  to  $10.4202 \pm 0.2230$ , increasing in rainy season and varies from  $7.1200 \pm 0.1254$  to  $9.1010 \pm 0.1022$ , whereas it is minimum in winter season and varies from  $4.1211 \pm 0.4212$  to  $5.270 \pm 0.3201$ .
  - **Gill glycogen:** The seasonal changes in the glycogen content in gill of *L. corrianus* are shown in (Table No. 5 and Figure No. 5). The percentage of glycogen was found to be maximum in summer and varies from  $7.1220 \pm 0.2728$  to  $10.2130 \pm 0.6302$ , increasing in rainy season and varies from  $6.5210 \pm 0.5312$  to  $9.2023 \pm 0.2112$ , whereas it is minimum in winter season and varies from  $4.2210 \pm 0.2120$  to  $5.1202 \pm 0.4226$ .

The present study revealed that, there is significant variation in the biochemical composition in different body tissues according to seasonal changes. Seasonal changes in protein content may be of great importance in relation to energy metabolism necessary for growth and reproduction (Lodeiros *et al.*, 2001). Organic constituents like protein act as key substances for different metabolic activities. Protein is the main organic nutrient used to build up different body tissues. All the tissues show increasing order protein contents in rainy season, which is correlated with highest body activities of animal during this season. And due to increase inflow and turbidity of water and to cope up with new environmental change. It might be due to favourable environmental lots of food availability and the period of growth with the gonadal development. Similar conclusions were reported in *M. edulis*, in British water by Williams, 1969 and Mane and Nagabhushanam, 1978.

The protein seems to be its only alternative resource of energy under conditions of food scarcity. During May 1st and 2nd fortnight the drastic environmental condition results in recovery of gonad tissue. Protein content decrease in gonad and hepatopancreas, during this period was seen in *L. corrianus* from Godavari River by (Muly, 1988). Thus, food availability may be the important source of nutrients required for the gonadal repining process. Seasonal variation in temperature and availability of food appear to be closely related to energy available for growth and reproduction in other bivalve species (Smaal *et al.*, 1997). In *E. exalbida* from Ushuaia Bay, shell growth in spring (Lomovasky *et al.*, 2002). Whereas the protein content showed a significant increase in Winter February 2001, however, it might be due to favorable environmental condition, lots of food availability and the period of growth with the gonadal development. The relative content of protein varies seasonally. These changes are principally related to the reproductive cycle and the season maximum shell growth.

Similar characteristics have been observed in other bivalve *Lyropecten (Nodipecten) nodosus* (Lodeiros *et al.*, 2001). The protein seems to be its only alternative resource of energy under conditions of food scarcity. However, it cannot be certain without further studies and proper investigation about the possible advantage of using protein as an energy reserve and the mechanisms of regulation (e.g., anti-freezing proteins). In Summer May 2000 the protein content was significantly low due to drastic environmental condition the rise in temperature, scarcity of food availability, starvation effect and endogenous role of hormone as the removal of cerebral ganglion maybe responsible of decrease in protein content. Glycogen estimation observed during experimental work has been given in Fig for *Lamellidens Corrianus*. The present study revealed that, there is significant variation in the biochemical composition in different body tissues according to seasonal changes. Seasonal changes in protein content may be of great importance in relation to energy metabolism necessary for growth and reproduction (Lodeiros *et al.*, 2001). Organic constituents like glycogen act as key substances for different metabolic activities. All the tissues show increasing order glycogen contents in rainy season, which is correlated with highest body activities of animal during this season. And due to increase inflow and turbidity of water and to cope up with new environmental change. It might be due to favourable environmental lots of food availability and the period of growth with the gonadal development. Similar conclusions were reported in *M. edulis*, in British water by Williams, 1969 and Mane and Nagabhushanam, 1978. Bivalves generally store carbohydrates in large amounts during their growing season and use them over the rest of the year (Beukema, 1997);

Glycogen is the primary energy store in bivalves (Garbbott, 1983). In the entire body organ it is observed that glycogen acontent is significantly accumulated is found to be more during summer season. The relationship of the energy transfer between different tissues, their capacity of reserve amounts under food availability, and their positive relationship with the high temperature and gonadal maturation have been shown in different species of bivalve molluscs such as scallops (Urrutia *et al.*, 2001). The relative content of glycogen vary seasonally. These changes are principally related to the reproductive cycle and the season maximum shell growth. Similar characteristics have been observed in other bivalve *Lyropecten (Nodipecten) nodosus* (Lodeiros *et al.*, 2001).

## REFERENCES

- Smaal, A. C., Vonck, A. P. M. A. and Bakker, M. 1997. Seasonal variation in physiological energetics of *Mytilus edulis* and *Cerastoderma edule* of different size classes. *J. Mar. Biol. Assoc. U. K.* 77: 817-838.
- Kapil Manoj and Ragothaman, G (1999) : Mercury, copper and cadmium induced changes in the proteins levels in muscle tissue of an edible estuarine fish, *Bolephthalmus dussumeri* (Cuv). *J. Environ. Biol.* 20(3): 231-234
- Lehinger A L 1984 In: Biochemistry 3<sup>rd</sup> ed., Kalyani Publisher, Ludhiana, New Delhi.
- Lodeiros, C. J., Rengel, J. J., Guderley, H. E., Nuseni, O. and Himmelman, J. H. 2001. Biochemical composition and energy allocation in the tropical scallop *Lyropecten (Nodipecten) nodosus* during the months leading up to and

- following the development gonads. *Aquaculture*. 199: 63-72.
- Lomovasky, B. J., Brey, T., Morriconi, E. and Calvo, J. 2002. Growth and production of the venerid bivalve *Eurhomalea exalbida* in the Beagle Channel, Tierra del Fuego. *J. Sea Res.* 48: 209-216
- Lowry O.H., Rosenbrough N.J., Farr A.L., and Randall R.J. 1951. Protein measurement with Folin phenol reagent. *J. Biol. Chem.* 193, 265-275.
- Muly, S. D. 1988. Reproductive physiology of Lamellibranches molluscs from Marathwada state, Ph. D Thesis, Marathwada University, Aurangabad.
- Nagabhushanam, R. and Mane, U. H. 1978. Seasonal variations in the biochemical composition of *Mytilus viridis* at Ratnagiri on the West Coast of India. *Hydrobiol.* 57: 69 – 72.
- Genton, Laurence; Melzer, Katarina; Pichard, Claude (2010). "Energy and macronutrient requirements for physical fitness in exercising subjects". *Clinical Nutrition* 29 (4): 413–423
- Williams, C. S. 1969. The effect of *Mytilicola intestinalis* on the biochemical composition of mussel. *J. Mar. Biol. Assoc. U.K.* 49: 161-173.
- Beukema, J. J. and De Bruin, W. 1977. Seasonal changes in dry weight and chemical composition of the soft parts of the tellinid bivalve *Macoma balthica* in the Dutch Wadden Sea. *Neth. J. Sea Res* 11(1), 42–55.
- Garbbott, P. A. 1983. Developmental and seasonal metabolic activities in marine molluscs. In the *Mollusca*. *Envi. Biochem. and Physiology*, P.W. Hochachka. Academic Press, New York. 2: 165-217.
- Lodeiros, C. J., Rengel, J. J., Guderley, H. E., Nuseni, O. and Himmelman, J. H. 2001. Biochemical composition and energy allocation in the tropical scallop *Lyropecten (Nodipecten) nodosus* during the months leading up to and following the development gonads. *Aquaculture*. 199: 63-72.
- Nagabhushanam, R. and Mane, U. H. 1978. Seasonal variations in the biochemical composition of *Mytilus viridis* at Ratnagiri on the West Coast of India. *Hydrobiol.* 57: 69 – 72.
- Prosser, C.L. (1973) "Comparative Animal Physiology" Edtd by C.L. Prosser, W.B. Sounder Co. Philadelphia, London, Toranto pp .317-335.
- Seifer S., Dayton S., Navie B., and Muntury G.R. 1950. The estimation of Glycogen with the anthrone reagent. *Arch. Biochem. Biophys.* 25 (1), 191-200
- Urrutia, G. X., Navarro, J. M., Closing, E. and Stead R. A. 2001. The effects of environmental factors on the biochemical composition of the bivalve *Tagelus dombeii* (Lamarck, 1818) (Tellinacea: Solccurtidae) from the intertidal flat of Coihuin, Puerto Montt, Chile. *J. Shellfish Res.* 20: 1077-1087
- Williams, C. S. 1969. The effect of *Mytilicola intestinalis* on the biochemical composition of mussel. *J. Mar. Biol. Assoc. U.K.* 49: 161-173.

\*\*\*\*\*