



ISSN: 0976-3376

Available Online at <http://www.journalajst.com>

ASIAN JOURNAL OF
SCIENCE AND TECHNOLOGY

Asian Journal of Science and Technology
Vol. 5, Issue 3, pp. 174-175, March, 2014

RESEARCH ARTICLE

EFFECT OF AEROBIC EXERCISE ON SELECTED PHYSIOLOGICAL VARIABLES AMONG OBESE MALE CHILDREN

*Prasana Sundara Raju, P.

Lecturer in Physical Education, Pondicherry Institute of Medical Science, Pondicherry

ARTICLE INFO

Article History:

Received 21st December, 2013

Received in revised form

13th January, 2014

Accepted 12th February, 2014

Published online 25th March, 2014

Key words:

Obese,

School children,

Aerobic capacity,

Percent body fat

ABSTRACT

The purpose of the study was to find out the effectiveness of aerobic training on selected physiological variables among children prone to obesity. To achieve the purpose twenty five schools boys were selected from Fathima Higher Secondary School, Karuvadikuppam, Puducherry and their age range between 14 to 16 years. The physiological variables confined to this study were maximum oxygen consumption and percent body fat were selected as criterion variables were assessed using Cooper's twelve minutes run/walk test and skin fold measurements, before and after the experimental treatment for six weeks. The data collected thereof on maximum oxygen consumption and percent body fat were statistically analysed using paired 't' ratio. The result of this study showed that significant improvement on maximal oxygen consumption and significant reduction on percent body fat. This study showed that six week of aerobic exercise significantly minimizes the cardiovascular risk among obese school children.

Copyright © 2014 Prasana Sundara Raju, P. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Obesity is recognized as a major global burden to health (Wearing *et al.*, 2006). In India when the child reaches adolescence their level of physical activity declines. There is evidence (Wang *et al.*, 2002) that children and adolescents of urban families are more overweight than rural, possibly because of decreased physical activities, sedentary lifestyle, altered eating patterns and increased fat content of the diet. Increase in sedentary activities, such as television viewing and computer games, is suspected to be responsible for the decline in physical activity levels. Obesity is associated with increased systemic blood pressure, decreased aerobic fitness, cardiopulmonary function, increased rate of Type 2 diabetes Mellitus and cardiovascular diseases (Wisloff *et al.*, 2007). In order to eradicate obesity it is therefore important to encourage sustainable physical activity habits in children, and further reinforcing these habits in children, which will help establish desirable healthy lifestyle patterns that continue into adulthood. The purpose of the study was to find out the effectiveness of aerobic training on selected physiological variables among children prone to obesity.

MATERIALS AND METHODS

Subjects

To achieve the purpose of the study, twenty five schools boys from Fathima Higher Secondary School, Karuvadikuppam,

Puducherry, in the age group of 14 to 16 years, with a body mass index (BMI) between 85th and 95th percentile based on the cut-off points of the CDC dataset for BMI were considered as overweight and selected as subjects for this study. The selected subjects neither have the experience of organised fitness training nor participating in any other special coaching programme. A qualified physician examined the subjects medically and declared that they were fit to undergo the physical activity programme. The subjects were free to withdraw their consent to participate in the training programme, in case they felt any discomfort during the period of training. But there were no dropouts in the study.

Variable and test

The experimental variable used in the present study was aerobic training for six weeks. The criterion variables chosen for the present research were physiological variables. Maximal oxygen consumption – Coopers 12 minutes run and walk test and percent body fat – skin fold caliper.

Training Programme

During the training period, the selected subjects underwent aerobic training programme for three days a week, for six weeks in addition to their regular school activities. The subjects underwent the training programme under strict supervision. Prior to every training session, the subjects underwent ten minutes of warming-up exercises, which included jogging, stretching and striding. The subjects

*Corresponding author: Prasana Sundara Raju, P., Lecturer in Physical Education, Pondicherry Institute of Medical Science, Pondicherry

involved in the training programme were questioned about their state throughout the training period, none of them reported any injuries, however muscle soreness was reported in the early weeks but it subsided later. Attendance was taken regularly. The training load was increased progressively as mentioned in the training schedule Table 1.

Table 1. Training Schedule

Week	Duration	Running pace	Distance
I Week	15 minutes	350 metres slow & 50 metres fast	2400
II Week	16 minutes	350 metres slow & 50 metres fast	2550
III Week	16 minutes	325 metres slow & 75 metres fast	2800
IV Week	17 minutes	325 metres slow & 75 metres fast	3000
V Week	17 minutes	150 metres slow & 50 metres fast	3400
VI Week	18 minutes	150 metres slow & 50 metres fast	3600

Statistical Techniques

The data collected from the subjects prior to and after experimentation on selected physiological parameters and body composition, that is, maximum oxygen consumption and percent body fat were statistically examined for significant alterations because of aerobic training, if any, by applying the dependant 't' test with the help of SPSS package. In determining the significance of 't' ratio the confidence interval was fixed at 0.05 level, which is considered appropriate enough for the study.

RESULTS

The analysis of data on physiological variables between pre test and post test capabilities were statistically analysed using 't' ratio and presented in Table 2.

Table 2. Computation of Data on Maximum Oxygen Consumption and Percent body fat

Variables	Test	Mean	SD	DM	Std Error of DM	't' ratio
Maximum Oxygen Consumption	Pre-test	23.928	2.04199	0.9032	0.04373	20.652*
	Post test	24.831	2.09928			
Percent body fat	Pre-test	25.76	1.00125	0.40	.0323	12.394*
	Post test	25.36	1.01571			

The required table value for significance at 0.05 level of confidence for df of 24 is 2.064. Table – 2 indicates that the pre test and post test means on maximum oxygen consumption were 23.928 and 24.831 respectively. The obtained 't' ratio of 20.652 on maximum oxygen consumption was greater than the required table value 2.064 for significance with df of 24 at 0.05 level of confidence. The result of the study showed that

there was significant improvement on maximum oxygen consumption. Similarly, the obtained 't' ratio of 12.394 on percent body fat was greater than the required table value 2.064 for significance with df of 24 at 0.05 level of confidence. The result of the study showed that there was a significant reduction on percent body fat.

DISCUSSION ON FINDINGS

The results obtained from this study on effectiveness of aerobic training on selected physiological parameters and body composition among children prone to obesity, clearly shows that there was a significant improvement on maximum oxygen consumption and a considerable reduction on percent body fat as a result of aerobic training programme for the period of six weeks. These results were also in line with the previous literature that found improvements in health related parameters of obese participants as a result of regular exercise participation (Steinbeck, 2001; Karacabey, 2009). Moreover, similar results have been reported in adults (Dengel *et al.*, 1998; Wong *et al.*, 2008).

Conclusion

It is concluded that aerobic exercise for six week significantly minimized the cardiovascular risk of obese children.

REFERENCES

- Dengel, D.R., Galecki, A.T., Hagberg, J.M. 1998. The independent and combined effects of weight loss and aerobic exercise on blood pressure and oral glucose tolerance in older men. *Am. J. Hypertens*, 11: 1405-12.
- Karacabey K. 2009. The effect of exercise on leptin, insulin, cortisol and lipid profiles in obese children. *J. Int. Med. Res*, 37(5): 1472- 1478.
- Steinbeck K.S. 2001. The importance of physical activity in the prevention of overweight and obesity in childhood: a review and an opinion. *Obesity Rev*, 2: 117-130.
- Wang Y, Monteiro C, Popkin BM. 2002. Trends of obesity and underweight in older children and adolescents in the United States, Brazil, China, and Russia. *Am J Clin Nutr*, 75:971–7.
- Wearing, *et al.* 2006. The impact of childhood obesity on musculoskeletal form. *Obesity reviews*, 7: 209-218
- Wisløff *et al.*, 2007. Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients: a randomized study. *Circulation*, 115, 3086–3094
- Wong *et al.*, 2008. Effects of a 12 week exercise training program on aerobic fitness, body composition, blood lipids and C-reactive protein in adolescents with obese. *Ann. Acad. Med. Singapore*, 37(4): 286-93.
